



Securing Eating Disorders Treatment: Ammunition for Arguments with Third Parties

Compiled and Written by Margo Maine, PhD

Argument # 1: A full course of treatment is cost-effective for eating disorders.

Crow and Nyman (2004)

Relative to other accepted medical interventions, the treatment of eating disorders has been shown to be cost-effective and, in fact, quite reasonable.

Halmi et al (2000)

Between 1985 & 1998 readmissions of eating disorder patients increased steadily as length of stay became briefer and weight at discharge was lower.

Commerford, Licino, and Halmi (1997)

Shorter periods of treatment for eating disorders are associated with less successful outcomes.

Halmi and Licino (1989)

Patients with anorexia nervosa who reached 98% of Ideal Body Weight (IBW) prior to discharge from inpatient are less likely to relapse than those who only achieved 83% IBW. (Program completion = abstinence from bingeing and purging and maintaining weight for 2 weeks.)

Baran et al (1995)

Compared 14 patients with anorexia nervosa who achieved normal weight (96% IBW) to 8 patients who reached only 76% IBW. (see below)

	<u>96%</u>	<u>76%</u>
Length of stay	116 days	46 days
Menstrual disturbances	21%	62%
Re-hospitalization	7%	62%
Persistent anorexic symptoms	19%	57%

Concluded that achieving IBW has better clinical course and may be more cost-effective in long run, especially in light of costs of medical problems (osteoporosis, infertility, dehydration, electrolyte imbalances, cardiac and other compromises).

Argument # 2: Specialized treatment for eating disorders is preferable and cost-effective.

Crisp, Callender, Halek and Hsu (1992)

Specialized treatment reduces mortality.

Argument # 3: Recovery takes place over a long period of time.

Strober, Freeman, Morrell (1997)

“The course of anorexia nervosa is protracted.” (p. 339)

76% of sample studied for 10-15 years after admission met criteria for full recovery, but time to recovery ranged from 57-74 months; 10% met criteria for partial recovery.



Ammunition for Arguments with Third Parties

Compiled and Written by Margo Maine, PhD

Argument # 4: Treatment of bulimia is effective.

Reas et al (2000)

Patients with bulimia nervosa demonstrate a better recovery rate if they receive treatment early in their illness.

If treated within the first 5 years, the recovery rate is 80%.

If not treated till after 15 years of symptoms, recovery falls to 20%.

Argument # 5: Even successful treatment has an uneven course.

Fichter and Quadflieg (1997)

Six year follow-up of 196 female bulimic patients indicate 59.9% had achieved a good outcome; 29.9% poor, and 1.1% deceased.

The course of recovery is uneven, with a decline during the 2 years after intensive inpatient treatment, then later improvement and stabilization.

Argument # 6: Eating disorders are serious and lethal.

Crow and Nyman (2004)

Long-term studies of anorexia nervosa report a mortality rate of 10%.

Herzog et al (2000)

The mortality rate for women suffering from eating disorders is 9.6 times the expected rate and the suicide rate is 58 times greater.

Sullivan (1995)

Conducted meta-analysis from 42 studies of patient mortality, finding 178 deaths in 3,006 patients:

54% from complications of the eating disorders

27% suicide

19% other/unknown

Rate of .56% mortality/years is 12 times greater than general death rate for women aged 15-24 and suicide rate is 75 times greater.

Argument # 7: The mortality rate increases with the duration of symptoms.

APA (2000)

The mortality rate at 5 years is 5%, increasing to 20% at 20 year follow-up.



Ammunition for Arguments with Third Parties

Compiled and Written by Margo Maine, PhD

Argument # 8: Comprehensive and long-term treatment does “pay off.”

Garfinkel (1995)

Outcome studies following patients for 5-10 years indicate mortality rate of 5%. Studies of patients followed for 20-30 years find 18% mortality rate from anorexia nervosa or suicide.

Treatment works. Of those followed 5-10 years, approximately half of sample had recovered, 25% improved with some residual symptoms, and 25% remain ill or die.

Argument # 9: Younger patients require intense and aggressive treatment.

Society for Adolescent Medicine (2003)

“Because of the potentially irreversible effects of an eating disorder on physical and emotional growth and development in adolescents, the high mortality and the evidence suggesting improved outcome with early treatment, the threshold for intervention in adolescents should be lower than in adults.” (p.500)

Irreversible risks are growth retardation, pubertal delay or arrest, impaired acquisition of peak bone mass, increased risk of osteoporosis.

Argument # 10: Utilization of mental health benefits may offset high medical costs associated with eating disorders.

APA Practice Guidelines (2000 & 1993)

Reports these medical findings:

- Physical consequences of eating disorders include all serious sequelae of malnutrition, especially cardiovascular compromise.
- Prepubertal patients may have arrested sexual maturity and growth failure.
- Even those who “look and feel deceptively well,” with normal EKG’s may have cardiac irregularities, variations with pulse and blood pressure, and are at risk for sudden death.
- Prolonged amenorrhea (>6 months) may result in irreversible osteopenia and a high rate of fractures.
- Abnormal CT scans of brain are found in >50% of patients with anorexia nervosa.
- Bulimic behaviors may result in electrolyte, fluid and mineral imbalance; may be presenting cardiac risk; gastric irritation and bleeds; large bowel abnormalities; dental enamel erosion; peripheral muscle weakness, cardiomyopathy, hypometabolism. Despite normal weights a bulimic may present cardiac risk; gastric irritation and bleeds; large bowel abnormalities; dental enamel erosion; peripheral muscle weakness, cardiomyopathy, hypometabolism. Despite normal weights a bulimic may be starving and severely malnourished.



Ammunition for Arguments with Third Parties

Compiled and Written by Margo Maine, PhD

BIBLIOGRAPHY

American Psychiatric Association (APA) (1993). Practice Guideline for Eating Disorders. American Journal of Psychiatry, 150 (2), 212-228.

American Psychiatric Association (APA) (2000). Practice Guideline for the Treatment of Patients with Eating Disorders (Revision). American Journal of Psychiatry, 157 (1), January Supplement 1-39.

Baran, S.A., Weltzin, T.E., Kaye, W.H. (1995). Low Discharge Weight and Outcome in Anorexia Nervosa. American Journal of Psychiatry, 152, 1070-72.

Commerford, M.C.; Licino, J. & Halmi, K.A. (1997). Guidelines for Discharging Eating Disorder Inpatients. Eating Disorders, 6, 69-74.

Crisp, A.H., Callender, J.S., Halek, E., Hsu, K.G. (1992). Long-term Mortality in Anorexia Nervosa: A 20 Year Follow-up of the St. George's and Aberdeen Cohorts. British Journal of Psychiatry, 161, 104-7.

Crow S. J. and Nyman, J.A. (2004). The Cost-effectiveness of Anorexia Nervosa Treatment. International Journal of Eating Disorders, 35, 155-60.

Fichter, M.M. and Quadflieg, N. (1997). Six-Year Course of Bulimia Nervosa. International Journal of Eating Disorders, 22 (3), 361-84.

Garfinkel, P.E. (1995). Eating Disorders. In H.I. Kaplan and B.J. Sadock (eds). Comprehensive Textbook of Psychiatry VI. 1361-71.

Halmi, K., Wiseman, K., Sunday, S., Harris, W., Klapper, F. (2000). The Changing Epidemiology of Hospitalized Eating Disorder Patients. Paper presented at Academy of Eating Disorders Conference, NY, May 4-7, 2000.

Halmi, K., Licino, E. (1989). Outcome: Hospital Program for Eating Disorders, in C.M.E. Syllabus and Proceedings Summary. 142nd Annual meeting of American Psychiatric Association, Washington, DC.

Herzog, D.B., Greenwood, D.N., Dorer, D.J., Flores, A.T., Ekeblad, E.R., Richards, A., Blais, M.A., Keller, M.B. (2000). Mortality in Eating Disorders: A Descriptive Study. International Journal of Eating Disorders, 28, 20-26.

Mehler, P.S., Anderson, A.E. (1999). Eating Disorders: A Guide to Medical Care & Complications. Baltimore, MD: Johns Hopkins University Press.

Reas, D.L., Williamson, D.A., Martin, C.K., Zucker, N.L. (2000). Duration of Illness Predicts Outcome for Bulimia Nervosa: A Long-term Follow-up Study. International Journal of Eating Disorders, 27 (4), 428-34.

Society for Adolescent Medicine (2003). Eating Disorders in Adolescents: Position Paper of the Society for Adolescent Medicine. Journal of Adolescent Health, 33, 496-503.

Strober, M., Freeman, R., and Morrell, W. (1997). The Long-Term, Course of Severe Anorexia Nervosa in Adolescents: Survival Analysis of Recovery, Relapse, and Outcome Predictors over 10-15 Years in a Prospective Study. International Journal of Eating Disorders, 22 (4), 339-60.

Sullivan, P.F. (1995). Mortality in Anorexia Nervosa. American Journal of Psychiatry, 152, 1073-74.

